



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.


We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Solution by the **PROPOSER**.

Let R = the radius of desert, T = time, v = rate. Let P be the position of the man at any instant. Draw about P an infinitesimal circle, MSK . Call the



angle MPN , θ . Then $\theta = \cos^{-1} \frac{PN}{PM}$.

Now the rate at which the man must approach the circumference in order to be off in a given time is R/T . In an infinitely small time the distance will be $(R/T)dt$.

Also $PN = vdt$.

$$\therefore \theta = \cos^{-1} \left(\frac{R}{Tv} \right) \dots \dots \dots (1).$$

Now R , T , and v are positive. Therefore the value of θ defined by equation (1) has to do with an angle less than 90° .

Now if the man at each instant goes within the angle MPN , he will get off the desert in the given time. The chance that he will do this is

$$C = \frac{2\cos^{-1}[(R/Tv)]}{\pi} \dots \dots \dots (2).$$

Hence the required probability in given by (2).

If $R=0$, or $T=\infty$, or $v=\infty$, $C=1$. If $R=Tv$, $C=0$.

If $R > Tv$, C is impossible.

Let $R=1$, and $v=1$. To find the time which he must have at his disposal in order that he may have half a chance to get off the desert.

Clearly $T = \sqrt{2}$.

PROBLEMS FOR SOLUTION.

ARITHMETIC.

75. Proposed by J. A. CALDERHEAD, M. Sc., Professor of Mathematics in Curry University, Pittsburg, Pennsylvania.

If 24 men, in 15 days of 12 hours each, dig a trench 300 yards long, 5 yards wide, 6 feet deep for 540 five-cent loaves when flour is \$8 a barrel; what is flour worth a barrel when 45 men, working $5\frac{1}{2}$ days of 10 hours each, dig a trench 125 yards long, 5 yards wide, 8 feet deep for 320 four-cent loaves? *Solve by proportion.*

76. Proposed by E. W. MORRELL, Professor of Mathematics in Montpelier Seminary, Montpelier, Vermont.

An eastern nobleman willed his entire estate to his three sons on the condition that the oldest should have one-half, the next one-third, and the youngest one-ninth. His estate, on inventory, was found to consist of 17 elephants. What should be the share of each?